

If you are an adult with nasal polyps who has had previous ethmoid sinus surgery,

## Take the fight to the source with the SINUVA® Sinus Implant



### SINUVA is an alternative to nasal surgery that



- Is placed during a routine office visit, with the entire procedure usually taking 30-40 minutes from start to finish
- Delivers medicine to shrink nasal polyps for up to 90 days
  - Reduces symptoms of nasal obstruction and congestion
  - Has been shown to improve a patient's altered sense of smell
- Is not for people with hypersensitivity to corticosteroids or those with nasal ulcers or trauma
- Has risks similar to those associated with other endoscopic sinus procedures, such as nose bleed, injuries to blood vessels, bacterial infection, irritation, or perforation



**To learn more, talk to your doctor or visit [www.SINUVA.com](http://www.SINUVA.com)**

SINUVA® Sinus Implant is a steroid-releasing (mometasone furoate) implant indicated for the treatment of nasal polyps in patients  $\geq 18$  years of age who have had ethmoid sinus surgery.

**Please see reverse for Important Safety Information about SINUVA and the accompanying Full Prescribing Information.**

**SINUVA®**  
(mometasone furoate) sinus implant



## INDICATION

SINUVA Sinus Implant is a prescription steroid-releasing (mometasone furoate) implant indicated for the treatment of nasal polyps in patients 18 years or older who have had ethmoid sinus surgery.

## IMPORTANT SAFETY INFORMATION

### Who should not use SINUVA?

Do not use SINUVA if you are allergic to mometasone furoate or any ingredients of the implant.

### What should I tell my doctor before receiving SINUVA?

Before you receive SINUVA, tell your doctor about all medical conditions you have including nasal/sinus problems (such as nasal ulcers or trauma), eye problems (such as glaucoma or cataracts), or any untreated fungal, bacterial, or viral infections.

### What are the possible side effects of SINUVA?

#### Serious side effects of SINUVA can include:

- **Local reactions** including nosebleed and injury to nerves or blood vessels in the nose/sinus.
- **Serious allergic reactions** have happened in patients using mometasone furoate including rash, itching or swelling of the lips, face, tongue, and throat, and breathing problems. Call your doctor right away if you have any of these reactions.
- **Weakened immune system** that may increase your risk of infections. Avoid contact with people who have contagious diseases such as chickenpox or measles. Call your doctor right away if you have been near someone with chickenpox or measles.
- **Adrenal insufficiency** is a condition in which the adrenal glands do not make enough steroid hormones and can cause tiredness, weakness, nausea and vomiting and low blood pressure. Talk to your doctor if steroid effects such as Cushing Syndrome and adrenal suppression appear.

**The most common side effects** of SINUVA in clinical studies were bronchitis, cold symptoms, middle ear infections, headache, lightheadedness or dizziness, asthma, and nosebleeds. The following adverse reactions have been identified during post-approval use of the SINUVA sinus implant. These events include implant migration, lack of efficacy, nasal pain, headache, and nosebleeds.

Tell your doctor if you have any side effects that bother you or don't go away.

Risks related with the insertion and removal of SINUVA are similar to other endoscopic sinus procedures.

SINUVA is made from materials designed to soften over time and may fall out of the nose on its own as polyps decrease or if you sneeze or blow your nose forcefully. The implant will be removed 90 days after placement or earlier at your doctor's discretion.

**Contact your doctor immediately** if you have any changes in vision, excessive nasal bleeding, symptoms of infection or symptoms suggesting that the implant has moved, such as irritation or a choking sensation in the back of the throat.

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.**

**For important risk and use information, please see Full Prescribing Information for SINUVA at [www.SINUVA.com/PI](http://www.SINUVA.com/PI)**



To learn more, talk to your doctor  
or visit [www.SINUVA.com](http://www.SINUVA.com)